

 Kilgen® Sterile eye drops Moxifloxacin 0.5%	SIDE EFFECTS: The most frequently reported ocular adverse effects are conjunctivitis, decreased visual acuity, dry eyes, keratitis, ocular discomfort, ocular hyperemia, ocular pain, ocular pruritus, subconjunctival haemorrhage and tearing.
COMPOSITION: Each ml of eye drops contains Moxifloxacin Hydrochloride INN equivalent to Moxifloxacin 5 mg.	
DESCRIPTION: The bacterial action of Moxifloxacin results from the interference with topoisomerase II and IV (DNA gyrase). Topoisomerases are essential enzymes that control DNA topology and assist in DNA replication, repair and transcription. It is effective against Beta-lactam and macrolide resistant bacteria.	USE IN PREGNANCY AND LACTATION: Since there are no adequate and well-controlled studies in pregnant women, this eye drops should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Moxifloxacin has not been measured in human; although it can be presumed to be excreted in human milk. Caution should be exercised when the eye drops is administered to a nursing mother.
INDICATIONS: Kilgen eye drops is indicated for the treatment of bacterial conjunctivitis caused by susceptible strains of the following organisms :	PAEDIATRIC USE: Safety and effectiveness in paediatric patients below the age of 1 year have not been established.
Aerobic Gram-positive micro-organisms: <i>Corynebacterium species</i> , <i>Micrococcus luteus</i> , <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Staphylococcus haemolyticus</i> , <i>Staphylococcus hominis</i> , <i>Staphylococcus warneri</i> , <i>Streptococcus pneumoniae</i> , <i>Streptococcus viridans group</i> .	WARNINGS: Not for injection. In patients receiving systemically administered Quinolones, including Moxifloxacin, serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported. After a few time some patients accompanied by some reactions i.e cardiovascular collapse, loss of consciousness, airway obstruction, dyspnea, urticaria and itching.
Aerobic Gram-negative microorganisms: <i>Haemophilus para influenzae</i> , <i>Haemophilus influenzae</i> . Other microorganisms: <i>Chlamydia trachomatis</i> .	PRECAUTIONS: As with other anti-infectives, prolonged use may result in over growth of non-susceptible organisms including fungi. If superinfection occurs, discontinue use and institute alternate therapy.
DOSAGE AND ADMINISTRATION: One drop in the infected eye (s) 3 times a day for 7 days.	PACKING: Each plastic dropper bottle contains 5 ml sterile eye drops.
CONTRAINDICATIONS: Kilgen eye drops is contraindicated in patients with a history of hypersensitivity to Moxifloxacin, to other Quinolones, or any of the components in this medication.	<i>Manufactured by:</i> Gaco Pharmaceuticals (G. A. Company Ltd.) Dhaka, Bangladesh